arXiv:astro-ph/0110606v1 29 Oct 2001

Gravitational Wave Induced Rotation of the Plane of Polarization of Pulsar Signals

A.R. Prasanna and S. Mohanty

Physical Research Laboratory

Navrangpura, Ahmedabad 380 009

India

Abstract

We derive in this an expression for the rotation of plane of polarization, of an electromagnetic wave, induced by the field of a gravitational wave propagating along the same direction $\approx \frac{G\mu d^2\Omega^4}{3\omega}$, ω and Ω being their respective frequencies using the geometrical optics limit of the wave equation for fields. Estimating the effect for the case of pulses from binary pulsars, one finds it too small to be observable, there could be other sources like inspiralling binary or an asymmetric neutron star where the effect could be in the observable region.

04.30.Nk , 42.15.i, 97.60Gb, 97.80d

The existence of gravitational waves is although confirmed from the binary pulsar data, attempts to detect directly have been pursued for over three decades with all the available technology. In this letter we pose a question whether one could detect these waves through their interaction with electromagnetic waves. In fact, both electromagnetism and gravitation being long range interactions, described through respective fields, it is possible that there exist an interaction between the two and some imprint of gravitational waves gets etched on electromagnetic waves through space time curvature. As it is far more easier to detect electromagnetic waves, we propose that using wave equation for electromagnetic field components on curved space time, one could get a dispersion relation in the geometrical optics limit that shows a formal rotation of the plane of polarization of the electromagnetic wave. The magnitude of the angle of rotation $\left(\sim \frac{GM\Omega^4}{kC^5}\right)$, which depends on the fourth power of the gravitational wave frequency could become detectable through pulsar radiation (Han et al 1998), for systems with very high frequency Ω.

It has been shown (Mohanty and Prasanna, 1998) that the wave equation for the electromagnetic field tensor in curved spacetime involves a coupling with the Riemann as well as Ricci curvatures of the background spacetime, and is given by

$$\nabla^\mu \nabla_\mu F_{\alpha\beta} + R_{\alpha\beta\mu\nu} F^{\mu\nu} + R^\mu{}_\alpha F_{\beta\mu} - R^\mu{}_\beta F_{\alpha\mu} = 0. \tag{1}$$

Considering the space time outside material distribution $R_{\mu\nu} = 0$, but having non-zero curvature produced by plane gravitational waves we look for a solution of (1) in the geometrical optics limit (eikonal approximation) with the ansatz

$$\begin{aligned} F_{\alpha\beta} &= e^{is/\epsilon} f_{\alpha\beta} \\ f_{\alpha\beta} &= \textstyle\sum_n \left(\frac{\epsilon}{i}\right)^n f_{n\ \alpha\beta} \end{aligned} \tag{2}$$

and

$$g_{\alpha\beta} = \eta_{\alpha\beta} + \hat{g}_{\alpha\beta} \tag{3}$$

$$\begin{aligned} \hat{g}_{\alpha\beta} &= e^{is/\epsilon} \gamma_{\alpha\beta} \\ \gamma_{\alpha\beta} &= \textstyle\sum_n \left(\frac{\epsilon}{i}\right)^n \gamma_{n\ \alpha\beta} \end{aligned} \tag{4}$$

(Ehlers and Prasanna, 1996), wherein the phase functions s (electromagnetic) and S (gravitational) are rapidly varying as compared to their respective amplitudes and the smallness parameter $\epsilon = \frac{\lambda}{L}$, is the ratio of wavelength to the background scale length. We shall first express eq. (1) in geodesic normal coordinates to get

$$\partial^\mu \partial_\mu\ F_{\alpha\beta} + \frac{1}{2}\left(g_{\beta\mu,\alpha\nu} + g_{\alpha\nu,\beta\mu} - g_{\alpha\mu,\beta\nu} - g_{\beta\nu,\alpha\mu}\right) F^{\mu\nu} = 0 \tag{5}$$

Using (2)-(4) in (5) and denoting the wave vectors

$$\partial_\mu s = k_\mu \text{ and } \partial_\mu S = \ell_\mu \tag{6}$$

one gets

$$\begin{aligned} & e^{is/\epsilon}\left\{-\tfrac{1}{\epsilon^2} k^\mu k_\mu f_{\alpha\beta} + \tfrac{i}{\epsilon}\left(\partial^\mu k_\mu + 2k_\mu \partial^\mu\right) f_{\alpha\beta} + \partial^\mu \partial_\mu f_{\alpha\beta}\right\} \\ & +\tfrac{1}{2} e^{iS/\epsilon}\left\{-\tfrac{1}{\epsilon^2}\left(\ell_\nu \ell_\alpha \gamma_{\beta\mu} + \ell_\beta \ell_\mu \gamma_{\nu\alpha} - \ell_\beta \ell_\nu \gamma_{\alpha\mu} - \ell_\alpha \ell_\mu \gamma_{\beta\nu}\right)\right. \\ & +\tfrac{i}{\epsilon}\left[\left(\partial_\nu \ell_\alpha \partial_\nu + \ell_\nu \partial_\alpha\right)\gamma_{\beta\mu} + \left(\partial_\mu \ell_\beta + \ell_\beta \partial_\mu + \ell_\mu \partial_\beta\right)\gamma_{\alpha\nu}\right. \\ & \left. -\left(\partial_\nu \ell_\beta + \ell_\beta \partial_\nu \partial_\beta\right)\gamma_{\alpha\mu} - \left(\partial_\nu \ell_\alpha + \ell_\alpha \partial_\mu + \ell_\mu \partial_\alpha\right)\gamma_{\beta\nu}\right] \\ & \left. +\left(\partial_\nu \partial_\alpha \gamma_{\beta\mu} + \ell_\beta \partial_\mu \gamma_{\nu\alpha} - \partial_\nu \partial_\beta \gamma_{\alpha\mu} - \partial_\mu \partial_\alpha \gamma_{\nu\beta}\right)\right\} e^{is/\epsilon} f^{\mu\nu} = 0 \end{aligned} \tag{7}$$

Collecting the zeroth order terms in the amplitudes the equation reduces to

$$k^\mu k_\mu f_{0\alpha\beta} + \frac{1}{2}\left\{\ell_\nu \ell_\alpha \gamma_{0\beta\mu} + \ell_\beta \ell_\mu \gamma_{\nu\alpha} - \ell_\beta \ell_\nu \gamma_{0\alpha\mu} - \ell_\alpha \ell_\mu \gamma_{0\beta\nu}\right\} f_0^{\mu\nu} = 0 \tag{8}$$

Assuming the curvature to be due to plane monochromatic gravitational wave propagating in the z-direction, the space time metric would be given by

$$dS^2 = dt^2 - (1 - h_{xx})\, dx^2 - (1 + h_{xx})\, dy^2 + 2h_{xy} dx dy - dz^2 \tag{9}$$

Further, considering the electromagnetic wave also to be along the z-direction, one has the folowing wave vectors

$$\begin{aligned} \ell_\alpha &= (0, 0, \ell, \Omega) \\ k_\alpha &= (0, 0, k, \omega) \end{aligned} \tag{10}$$

for the gravitational and electromagnetic waves respectively, and equation (7) takes the form:

$$k^\mu k_\mu f_{oi} + e^{iS/\epsilon} \Omega \left(\ell_\nu \gamma_{ij} f^{j\nu} \right) = 0 \tag{11}$$

which may be written for the electric fields f^{01} and f^{02}

$$\begin{pmatrix} (\omega^2 - k^2) + \Omega^2 \gamma_{11} e^{iS/\epsilon} & \Omega^2 \gamma_{12} e^{iS/\epsilon} \\ \Omega^2 \gamma_{21} e^{iS/\epsilon} & (\omega^2 - k^2) + \Omega^2 \gamma_{22} e^{iS/\epsilon} \end{pmatrix} \begin{pmatrix} f^{01} \\ f^{02} \end{pmatrix} = 0 \tag{12}$$

The condition for the non-trivial solution for f^{01}, f^{02} yields the dispersion relation

$$\left(\omega^2 - k^2 \right)^2 - \Omega^4 \left(\gamma_{11}^2 + \gamma_{12}^2 \right) e^{2iS/\epsilon} = 0$$

or

$$k^2 = \omega^2 \pm \Omega^2 \sqrt{\gamma_{11}^2 + \gamma_{12}^2} \;\; e^{iS/\epsilon} \tag{13}$$

It is well known that for the field of the weak gravitational wave, from the point masses in a Keplerian orbit (binary system), the metric potentials are given by (Peters and Mathews, 1963)

$$\begin{aligned} h_{xx} &= -\tfrac{G\Omega^2}{3r} \left(Q_{xx} - Q_{yy} \right) e^{i\Omega(t-z)} = -h_{yy} \\ h_{xy} &= -\tfrac{2G\Omega^2}{3r} Q_{xy} e^{i\Omega(t-z)} \end{aligned} \tag{14}$$

with

$$\begin{aligned} Q_{xx} &= \mu d^2 \cos^2 \psi \\ Q_{yy} &= \mu d^2 \sin^2 \psi \\ Q_{xy} &= \mu d^2 \sin \psi \cos \psi \end{aligned} \tag{15}$$

$\mu = \frac{M_1 M_2}{M_1 + M_2}$ being the reduced mass of M_1, M_2,the two pulsar masses and $d = a\frac{\left(1-e^2\right)}{(1+e\cos\psi)}$ while the angular velocity

$$\dot{\psi} = \frac{\left[G\left(M_1 + M_2\right)a\left(1-e\right)\right]^{1/2}}{d} \tag{16}$$

As we have $\gamma_{11} = h_{xx}e^{iS/\epsilon}$, $\gamma_{12} = h_{xy}e^{-iS/\epsilon}$, using (14) and (15) in (13), one finds

$$k^2 = \omega^2 \left(1 \pm \frac{G\mu d^2 \Omega^4}{3\omega^2 r} e^{i\Omega(t-z)}\right) \tag{17}$$

Since the gravitational waves propagate with the same phase velocity as electromagnetic waves, one gets the dispersion relation for electromagnetic waves to be

$$k_{\pm} = \omega \pm \frac{G\mu d^2 \Omega^4}{3\omega r} \tag{18}$$

Thus the electromagnetic pulse passing through the field of gravitational waves will have its plane of polarisation rotated through an angle

$$\Delta\phi = \int_{r_{min}}^{r_{max}} dr\left(k_+ - k_-\right) = \frac{G\mu d^2 \Omega^4}{3\omega} \ell n \left(\frac{r_{max}}{r_{min}}\right) \tag{19}$$

Consider the follwing examples:

(a) For the Hulse-Taylor binary 1913+16, the values of the relevant parameters are $\mu = 0.7 M_\odot$, $d \sim 2$ secs, orbital period $= P = \frac{2\pi}{\Omega} = 2.7906.98$ sec (Weisberg and Taylor 1984) . Considering the pulsar beam of frequency $\omega \approx 400$ MHz, the optical rotation over a distance of $\sim 10^2 kpc$ is $\Delta\phi \approx 10^{-28} rad$, which is very small.

(b) On the other hand, if one considers the gravitational waves from an inspiralling binary (Nakamura et al 1998) with a pulsar with $d \approx 100km$, between the two compact objects, the orbital frequency $\Omega = \left[\frac{G(M_1+M_2)}{d^3}\right]^{1/4} \sim 5 \times 10^{-3}$ sec. for $M_1 \sim M_2 \sim 1.4\ M_\odot$. For this system, the optical rotation of a 400 MHz radio signal over a distance of $\sim 100Mpc$ turns out to be $\Delta\phi \approx 1.9 \times 10^{-10} rad$.

(c) Finally considering the gravitational waves from an assymetric pulsar (Thorne 1997) with a non-zero quadrupole moment parametrized by ηGMR^2, the emitted gravitational wave will have the amplitude $h \approx \frac{\eta GMR^2\Omega^2}{r}$. Using this for the metric potential $h_{xx} = h_{yy} \approx h$, the dispersion relation for the electromagnetic signal would be

$$k_\pm = \omega\left[1 \pm \frac{\eta GMR^2\Omega^4}{\sqrt{2}r\omega^2}\right] \tag{20}$$

and the corresponding optical rotation, for $M \sim 1.4M_\odot$, $R \sim 10$ km, $\Omega = \frac{2\pi}{10^{-3}}s$, over a distance of 100 Mp_c is

$$\Delta\phi \approx 2.8\eta \times 10^{-8} \text{ radians} \tag{21}$$

If η is not too small, it may be possible to observe the signature of gravitational waves from such a pulsar.

In conclusion, it may be noted that the electromagnetic radiation emitted from a region which is also emitting sufficiently strong gravitational radiation, would suffer gravity induced birefringence which over a long distance can materialise as a rotation of the plane of polarization of the electromagnetic signal, thus establishing the presence of gravitational waves.

References

1. Ehlers J. and Prasanna A.R. (1996) *Class. & Quan. Grav.*, **13**, 2231.
2. Han J.L, Manchester R.N, Xu R.X , Qiao C.J,(1998) MNRAS **300**,373.
3. Mohanty S. and Prasanna A.R. (1998) *Nucl. Phys.* **B526**, 501.
4. Peters P.C. and Mathews J. (1963) *Phys. Rev.* **131**, 435.
5. Weisberg J.M and Taylor, J. H (1984), Phys Rev Lett **52**,1348.
6. Nakamura T, Sasaki M,Tanaka T,Thorne K.S, (1997) ApJ **487**,L139.
7. Thorne K.S, (1997) Archive gr-qc /9704042